arXiv:1208.6184v1 [math.CA] 30 Aug 2012

On a Local Mean Oscillation Decomposition

Jonathan Poelhuis and Alberto Torchinsky

August 31, 2012

The "local mean oscillation" decomposition of Lerner has proven particularly useful in recent literature. Such a functional decomposition was first proved for averages by Garnett and Jones [3], then suggested for medians by Fujii [2] and proved by Lerner [5], [6]. In this note we generate two local median oscillation decompositions using medians rather than rearrangements, so they apply to arbitrary measurable functions.

In what follows, we adopt the notations of [8] and [9]. In particular, for a cube $Q \subset \mathbb{R}^n$ and $0 < t < 1$, we say that $m_f(t, Q) = \sup\{M : |\{y \in Q : f(y) < M\}| \le t|Q|\}$ is the (maximal) median of f over Q with parameter t. For a cube $Q_0 \subset \mathbb{R}^n$ and $0 < s \le 1/2$, the local sharp maximal function restricted to Q_0 of a measurable function f at $x \in Q_0$ is

$$M^{\sharp}_{0,s,Q_0} f(x) = \sup_{x \in Q \subset Q_0} \inf_{c} \inf\{\alpha \ge 0 : |\{y \in Q : |f(y) - c| > \alpha\}| < s|Q|\}.$$

Additionally, for a cube $Q \subset \mathbb{R}^n$, let $\mathcal{D}(Q)$ denote the family of dyadic subcubes relative to Q; that is to say, those formed by repeated dyadic subdivision of Q into 2^n congruent subcubes. We will consider the maximal function

$$m^{t,\Delta}_{Q_0} g(x) = \sup_{Q \in \mathcal{D}(Q_0), x \in Q} |m_g(t, Q)|.$$

Finally, let $\widehat{Q}$ denote the dyadic parent of a cube Q.

The first local median oscillation decomposition holds for a range of indices, and so it extends Lerner's decomposition, which corresponds to the case $t = 1/2$, $s = 1/4$ in Theorem 1. On the other hand, the bound below is larger than his, but is sufficient for some applications such as those in [5].

Theorem 1. *Let f be a measurable function on a fixed cube $Q_0 \subset \mathbb{R}^n$, $0 < s < 1/2$, and $1/2 \le t < 1 - s$. Then there exists a (possibly empty) collection of subcubes $\{Q_j^v\} \subset \mathcal{D}(Q_0)$ and a family of collections $\{I_2^v\}_v$ such that*

(i) *For a.e. $x \in Q_0$,*

$$|f(x) - m_f(t, Q_0)| \le$$
$$4M^{\sharp}_{0,s,Q_0} f(x) + \sum_{v=1}^{\infty} \sum_{j \in I_2^v} \left(10n \inf_{y \in Q_j^v} M^{\sharp}_{0,s,\widehat{Q_j^v}} f(y) + 2 \inf_{y \in Q_j^v} M^{\sharp}_{0,s,Q_j^v} f(y) \right) \mathbb{1}_{Q_j^v}(x).$$

(ii) *For fixed v, $\{Q_j^v\}$ are nonoverlapping.*

(iii) *If* $\Omega^v = \bigcup_j Q_j^v$, *then* $\Omega^{v+1} \subset \Omega^v$.

(iv) *For all* j, $|\Omega^{v+1} \cap Q_j^v| \le \frac{s}{1-t}|Q_j^v|$.

Proof. Let $E^1 = \{x \in Q_0 : |f(x) - m_f(t, Q_0)| > 2\inf_{y\in Q_0} M^\sharp_{0,s,Q_0} f(y)\}$. If $|E^1| = 0$, the decomposition halts - trivially, for a.e. $x \in Q_0$,

$$|f(x) - m_f(t, Q_0)| \le 2 \inf_{y\in Q_0} M^\sharp_{0,s,Q_0} f(y). \tag{1}$$

So suppose that $|E^1| > 0$. Recall that by Lemma 4.1 in [8], for $\eta > 0$,

$$|\{x \in Q_0 : |f(x) - m_f(t, Q_0)| \ge 2 \inf_{y\in Q_0} M^\sharp_{0,s,Q_0} f(y) + \eta\}| < s|Q_0|.$$

Thus, picking $\eta_k \to 0^+$, by continuity from below it readily follows that

$$|\{x \in Q_0 : |f(x) - m_f(t, Q_0)| > 2 \inf_{y\in Q_0} M^\sharp_{0,s,Q_0} f(y)\}| \le s|Q_0|. \tag{2}$$

Now let $f^0 = (f - m_f(t, Q_0))\, \mathbb{1}_{Q_0}$, and

$$\Omega^1 = \{x \in Q_0 : m^{t,\Delta}_{Q_0}(f^0)(x) > 2 \inf_{y\in Q_0} M^\sharp_{0,s,Q_0} f(y)\}.$$

Then by Theorem 2.1 in [8], $E^1 \subset \Omega^1$ and $|\Omega^1| > 0$ as well. Write $\Omega^1 = \bigcup_j Q_j^1$ where the Q_j^1's are nonoverlapping maximal dyadic subcubes of Q_0 such that

$$|m_{f^0}(t, Q_j^1)| > 2 \inf_{y\in Q_0} M^\sharp_{0,s,Q_0} f(y),\ |m_{f^0}(t, \widehat{Q_j^1})| \le 2 \inf_{y\in Q_0} M^\sharp_{0,s,Q_0} f(y). \tag{3}$$

Since $m_{f^0}(t, Q_0) = 0$, $Q_j^1 \ne Q_0$ for any j.

Now since $t \ge 1/2$, from (1.10) in [8] it follows that

$$2 \inf_{y\in Q_0} M^\sharp_{0,s,Q_0} f(y) < |m_{f^0}(t, Q_j^1)| \le m_{|f^0|}(t, Q_j^1), \tag{4}$$

and therefore by the definition of median

$$|\{x \in Q_j^1 : |f^0(x)| > 2 \inf_{y\in Q_0} M^\sharp_{0,s,Q_0}\}| \ge (1-t)|Q_j^1|. \tag{5}$$

When these are summed, we have by (2) that

$$\begin{aligned}(1-t)\sum_j |Q_j^1| \le \sum_j &|\{x \in Q_j^1 : |f^0(x)| > 2 \inf_{y\in Q_0} M^\sharp_{0,s,Q_0} f(y)\}| \\ &\le |\{x \in Q_0 : |f^0(x)| > 2 \inf_{y\in Q_0} M^\sharp_{0,s,Q_0} f(y)\}| \le s|Q_0|,\end{aligned}$$

so that

$$\sum_j |Q_j^1| \le \frac{s}{1-t}|Q_0|, \tag{6}$$

where by the choice of s and t, $s/(1-t) < 1$.

Let $\alpha_j^1 = m_{f^0}(t, Q_j^1)$. By Lemma 4.3 in [8], we see that

$$|m_{f^0}(t, Q_j^1) - m_{f^0}(t, \widehat{Q_j^1})| \leq 10n \inf_{y \in Q_j^1} M^\sharp_{0,s,\widehat{Q_j^1}} f(y), \tag{7}$$

and therefore by (3) and (7)

$$\begin{aligned} |\alpha_j^1| &\leq |m_{f^0}(t, Q_j^1) - m_{f^0}(t, \widehat{Q_j^1})| + |m_{f_0}(t, \widehat{Q_j^1})| \\ &\leq 10n \inf_{y \in Q_j^1} M^\sharp_{0,s,\widehat{Q_j^1}} f(y) + 2 \inf_{y \in Q_0} M^\sharp_{0,s,Q_0} f(y). \end{aligned} \tag{8}$$

Then the first iteration of the local median oscillation decomposition of f when $|E^1| > 0$ is as follows: for a.e. $x \in Q_0$, with $g^1 = f^0 \mathbb{1}_{Q_0 \setminus \Omega^1}$,

$$f^0(x) = g^1(x) + \sum_j \alpha_j^1 \mathbb{1}_{Q_j^1}(x) + \sum_j \Big(f^0(x) - m_{f^0}(t, Q_j^1) \Big) \mathbb{1}_{Q_j^1}(x).$$

Note that g^1 has support off Ω^1, and clearly for a.e. $x \in Q_0$

$$|g^1(x)| \leq 2 \inf_{y \in Q_0} M^\sharp_{0,s,Q_0} f(y).$$

Now focus on the second sum. Since $f^0(x) - m_{f^0}(t, Q) = f(x) - m_f(t, Q)$ for all cubes Q and functions f supported in Q, this sum equals

$$\sum_j \left(f(x) - m_f(t, Q_j^1) \right) \mathbb{1}_{Q_j^1}(x).$$

The idea is to repeat the above argument for each function $f_j^1 = \Big(f - m_f(t, Q_j^1) \Big) \mathbb{1}_{Q_j^1}$, and so on.

We now describe the iteration. Assuming that $\{Q_j^{k-1}\}$ are the dyadic cubes corresponding to the $(k-1)$st generation of subcubes of Q_0 obtained as above, let

$$f_j^{k-1} = \Big(f - m_f(t, Q_j^{k-1}) \Big) \mathbb{1}_{Q_j^{k-1}},$$

and

$$E_j^k = \{ x \in Q_j^{k-1} : f_j^{k-1}(x) > 2 \inf_{y \in Q_j^{k-1}} M^\sharp_{0,s,Q_j^{k-1}} f(y) \}.$$

If $|E_j^k| = 0$, as in (1), we write $s_j^k = f_j^{k-1}$ which satisfies

$$|s_j^k(x)| \leq 2 \inf_{y \in Q_j^{k-1}} M^\sharp_{0,s,Q_j^{k-1}} f(y) \tag{9}$$

for a.e. $x \in Q_j^{k-1}$. These are the "s" functions since the decomposition "stops" at Q_j^{k-1}: clearly s_j^k has its support on Q_j^{k-1}, and Q_j^{k-1} contains no further subcubes of the decomposition.

If $|E_j^k| > 0$, we define

$$\Omega_j^k = \{ x \in Q_j^{k-1} : m^{t,\Delta}_{Q_j^{k-1}} f_j^{k-1}(x) > 2 \inf_{y \in Q_j^{k-1}} M^\sharp_{0,s,Q_j^{k-1}} f(y) \}.$$

Note that the E_j^k, and thus the Ω_j^k, are nonoverlapping. Then $|\Omega_j^k| > 0$ as well, and

$$\Omega_j^k = \bigcup_i Q_i^k,$$

where the Q_i^k's are nonoverlapping maximal dyadic subcubes of Q_j^{k-1} such that

$$|m_{f_j^{k-1}}(t, Q_i^k)| > 2 \inf_{y \in Q_j^{k-1}} M^\sharp_{0,s,Q_j^{k-1}} f(y), \; |m_{f_j^{k-1}}(t, \widehat{Q_i^k})| \le 2 \inf_{y \in Q_j^{k-1}} M^\sharp_{0,s,Q_j^{k-1}} f(y). \tag{10}$$

Then define

$$\Omega^k = \bigcup_j \Omega_j^k.$$

Let $\alpha_i^{k,j} = m_{f_j^{k-1}}(t, Q_i^k)$, and note that by (10)

$$\begin{aligned} |\alpha_i^{k,j}| &\le |m_{f_j^{k-1}}(t, Q_i^k) - m_{f_j^{k-1}}(t, \widehat{Q_i^k})| + |m_{f_j^{k-1}}(t, \widehat{Q_i^k})| \\ &\le 10n \inf_{y \in Q_i^k} M^\sharp_{0,s,\widehat{Q_i^k}} f(y) + 2 \inf_{y \in Q_j^{k-1}} M^\sharp_{0,s,Q_j^{k-1}} f(y). \end{aligned} \tag{11}$$

We then have

$$f_j^{k-1}(x) = g_j^k(x) + \sum_i \alpha_i^{k,j} \mathbb{1}_{Q_i^k}(x) + \sum_i \Big(f(x) - m_f(t, Q_i^k) \Big) \mathbb{1}_{Q_i^k}(x),$$

for a.e. $x \in Q_j^{k-1}$, where $g_j^k = f_j^{k-1} \mathbb{1}_{Q_j^{k-1} \setminus \Omega_j^k}$ clearly satisfies

$$|g_j^k(x)| \le 2 \inf_{y \in Q_j^{k-1}} M^\sharp_{0,s,Q_j^{k-1}} f(y) \tag{12}$$

for a.e. $x \in Q_j^{k-1}$. These are the "g" functions since the decomposition "goes on" or continues, into Q_j^{k-1}: g_j^k has support on Q_j^{k-1} away from Ω_j^k, which are the next subcubes in the decomposition.

We separate the Q_j^{k-1} into two families. One family, indexed by I_1^k, contains those cubes where the decomposition stops, and the other, indexed by I_2^k, where it continues. Specifically, let

$$I_1^k = \{j : \Omega^k \cap Q_j^{k-1} = \emptyset\}, \quad I_2^k = \{j : \Omega^k \cap Q_j^{k-1} \ne \emptyset\}.$$

Now we group the Q_i^k based on which Q_j^{k-1} contains them: if $j \in I_2^k$, let

$$J_j^k = \{i : Q_i^k \subset Q_j^{k-1}\}.$$

These definitions then give that

$$\Omega_j^k = \bigcup_{i \in J_j^k} Q_i^k.$$

Note that, as in (6),

$$|\Omega_j^k \cap Q_j^{k-1}| = \sum_{i \in J_j^k} |Q_i^k| \le \frac{s}{1-t} |Q_j^{k-1}|$$

so that

$$|\Omega^k| = \sum_j |\Omega_j^k \cap Q_j^{k-1}| \le \left(\frac{s}{1-t}\right) \sum_j |Q_j^{k-1}| = \left(\frac{s}{1-t}\right) |\Omega^{k-1}| \le \left(\frac{s}{1-t}\right)^k |Q_0|. \tag{13}$$

The kth iteration of the local median oscillation decomposition of the function f is as follows: for a.e. $x \in Q_0$,

$$f(x) - m_f(t, Q_0) = \sum_{v=1}^{k} \left(\sum_{j \in I_1^v} s_j^v + \sum_{j \in I_2^v} g_j^v \right) + \sum_{v=1}^{k} \sum_{j \in I_2^v} \sum_{i \in J_j^v} \alpha_i^{v,j} \mathbb{1}_{Q_i^v}(x) + \psi^k(x),$$

where

$$\psi^k = \sum_{j \in I_2^k} \sum_{i \in J_j^k} \left(f - m_f(t, Q_i^k) \right) \mathbb{1}_{Q_i^k}.$$

Since ψ^k is supported in Ω^k, by (13) it readily follows that $\psi^k \to 0$ a.e. in Q_0 as $k \to \infty$, and therefore

$$\begin{aligned} f(x) - m_f(t, Q_0) &= \sum_{v=1}^{\infty} \left(\sum_{j \in I_1^v} s_j^v + \sum_{j \in I_2^v} g_j^v \right) + \sum_{v=1}^{\infty} \sum_{j \in I_2^v} \sum_{i \in J_j^v} \alpha_i^{v,j} \mathbb{1}_{Q_j^v}(x) \\ &= S_1(x) + S_2(x), \end{aligned}$$

say.

In order to bound $|f(x) - m_f(t, Q_0)|$, consider first S_1. Of course, for all v and j the s_j^v's have nonoverlapping support. This is also true for the g_j^v's. Furthermore, the support of any g_j^v is nonoverlapping with that of any s_j^v. So for every v, j, and a.e. $x \in Q_0$, by (9) and (12)

$$\begin{aligned} \left| \sum_{v=1}^{\infty} \left(\sum_{j \in I_1^v} s_j^v + \sum_{j \in I_2^v} g_j^v \right) \right| &\le \max \left\{ \sup_{j \in I_1^v} \left\| f_j^{v-1} \right\|_{L^\infty}, \sup_{j \in I_2^v} \left\| f_j^{v-1} \mathbb{1}_{Q_j^{v-1} \setminus \Omega_j^v}(x) \right\|_{L^\infty} \right\} \\ &\le \max \left\{ \sup_{j \in I_1^v} \left(2 \inf_{y \in Q_j^{v-1}} M^\sharp_{0,s,Q_j^{v-1}} f(y) \right), \sup_{j \in I_2^v} \left(2 \inf_{y \in Q_j^{v-1}} M^\sharp_{0,s,Q_j^{v-1}} f(y) \right) \right\} \\ &\le 2 M^\sharp_{0,s,Q_0} f(x). \end{aligned} \tag{14}$$

Let's now consider S_2. The summand for $v = 1$ is distinguished, so we deal with it separately. By (8) above,

$$\begin{aligned} \left| \sum_j \alpha_j^1 \mathbb{1}_{Q_j^1}(x) \right| &\le \sum_j |\alpha_j^1| \mathbb{1}_{Q_j^1}(x) \\ &\le \sum_j \left(10n \inf_{y \in Q_j^1} M^\sharp_{0,s,\widehat{Q_j^1}} f(y) + 2 \inf_{y \in Q_0} M^\sharp_{0,s,Q_0} f(y) \right) \mathbb{1}_{Q_j^1}(x) \\ &\le \sum_j \left(10n \inf_{y \in Q_j^1} M^\sharp_{0,s,\widehat{Q_j^1}} f(y) \right) \mathbb{1}_{Q_j^1}(x) + 2 \inf_{y \in Q_0} M^\sharp_{0,s,Q_0} f(y). \end{aligned} \tag{15}$$

As for the other terms of the sum, by (11) we have

$$\begin{aligned}\Big|\sum_{v=2}^{\infty}\sum_{j\in I_2^v}\sum_{i\in J_j^v}\alpha_i^{v,j}\mathbb{1}_{Q_i^v}(x)\Big| &\le \sum_{v=2}^{\infty}\sum_{j\in I_2^v}\sum_{i\in J_j^v}|\alpha_i^{v,j}|\mathbb{1}_{Q_i^v}(x)\\ &\le \sum_{v=2}^{\infty}\sum_{j\in I_2^v}\sum_{i\in J_j^v}\Big(10n\inf_{y\in Q_i^v}M^\sharp_{0,s,\widehat{Q_i^v}}f(y)+2\inf_{y\in Q_j^{v-1}}M^\sharp_{0,s,Q_j^{v-1}}f(y)\Big)\mathbb{1}_{Q_i^v}(x)\\ &\le \sum_{v=2}^{\infty}\sum_{j\in I_2^v}\sum_{i\in J_j^v}\Big(10n\inf_{y\in Q_i^v}M^\sharp_{0,s,\widehat{Q_i^v}}f(y)\Big)\mathbb{1}_{Q_i^v}(x)\\ &\qquad +\sum_{v=2}^{\infty}\sum_{j\in I_2^v}\Big(2\inf_{y\in Q_j^{v-1}}M^\sharp_{0,s,Q_j^{v-1}}f(y)\Big)\mathbb{1}_{Q_j^{v-1}}(x). \end{aligned}\tag{16}$$

We combine (15) and (16) and note that since the sum is infinite and the families I_2^v are nested,

$$\begin{aligned}\Big|\sum_{v=1}^{\infty}\sum_{j\in I_2^v}\sum_{i\in J_j^v}\alpha_i^{v,j}\mathbb{1}_{Q_i^v}(x)\Big| &\le \sum_{v=1}^{\infty}\sum_{j\in I_2^v}\sum_{i\in J_j^v}\Big(10n\inf_{y\in Q_i^v}M^\sharp_{0,s,\widehat{Q_i^v}}f(y)\Big)\mathbb{1}_{Q_i^v}(x)\\ &\qquad +\sum_{v=1}^{\infty}\sum_{j\in I_2^v}\Big(2\inf_{y\in Q_j^{v-1}}M^\sharp_{0,s,Q_j^{v-1}}f(y)\Big)\mathbb{1}_{Q_j^{v-1}}(x)\\ &\le \sum_{v=1}^{\infty}\sum_{j\in I_2^v}\Big(10n\inf_{y\in Q_j^v}M^\sharp_{0,s,\widehat{Q_j^v}}f(y)+2\inf_{y\in Q_j^v}M^\sharp_{0,s,Q_j^v}f(y)\Big)\mathbb{1}_{Q_j^v}(x)\\ &\qquad +2\inf_{y\in Q_0}M^\sharp_{0,s,Q_0}f(y). \end{aligned}\tag{17}$$

Combining (14) and (17), finally we get that for a.e. $x\in Q_0$

$$\begin{aligned}&|f(x)-m_f(t,Q_0)|\le\\ &\qquad 4M^\sharp_{0,s,Q_0}f(x)+\sum_{v=1}^{\infty}\sum_{j\in I_2^v}\Big(10n\inf_{y\in Q_j^v}M^\sharp_{0,s,\widehat{Q_j^v}}f(y)+2\inf_{y\in Q_j^v}M^\sharp_{0,s,Q_j^v}f(y)\Big)\mathbb{1}_{Q_j^v}(x).\end{aligned}$$

Thus the conclusion holds.

□

The theorem and proof below are similar to those given in [5] and [6], but we present it to display how this decomposition leads to the same result. In what follows, M is the Hardy-Littlewood maximal function.

Theorem 2. *For any weight w, $0<s<1/2$, $1/2\le t<1-s$, $0<\delta\le 1$, and f measurable on a cube $Q_0\subset\mathbb{R}^n$,*

$$\int_{Q_0}|f(x)-m_f(t,Q_0)|w(x)dx\le c_{n,s,t}\int_{\mathbb{R}^n}\Big(M^\sharp_{0,s,Q_0}f(x)\Big)^{\delta}M\Big(\Big(M^\sharp_{0,s,Q_0}f(\cdot)\Big)^{1-\delta}w(\cdot)\Big)(x)dx.$$

If f is measurable on $\mathbb{R}^n$ such that $m_f(t,Q_0)\to 0$ as $Q_0\to\mathbb{R}^n$, then

$$\int_{\mathbb{R}^n}|f(x)|w(x)dx\le c_{n,s,t}\int_{\mathbb{R}^n}\Big(M^\sharp_{0,s}f(x)\Big)^\delta M\Big(\Big(M^\sharp_{0,s}f(\cdot)\Big)^{1-\delta}w(\cdot)\Big)(x)dx.$$

Proof. Applying Theorem 1, we have

$$\begin{aligned}\int_{Q_0}|f(x)-m_f(t,Q_0)|w(x)dx &\le 4\int_{Q_0}M^\sharp_{0,s,Q_0}f(x)\,w(x)dx\\ &\quad+\sum_{v=1}^{\infty}\sum_{j\in I_2^v}\left(10n\inf_{y\in Q_j^v}M^\sharp_{0,s,\widehat{Q_j^v}}f(y)+2\inf_{y\in Q_j^v}M^\sharp_{0,s,Q_j^v}f(y)\right)\int_{Q_j^v}w(x)dx.\end{aligned}$$

Now it easily follows that

$$\int_{Q_0}M^\sharp_{0,s,Q_0}f(x)\,w(x)dx\le\int_{\mathbb{R}^n}\Big(M^\sharp_{0,s,Q_0}f(x)\Big)^\delta M\Big(\Big(M^\sharp_{0,s,Q_0}f(\cdot)\Big)^{1-\delta}w(\cdot)\Big)(x)dx.$$

By construction, the Q_j^v are nonoverlapping over fixed v, but not over all v; indeed, the Q_j^v are subcubes of some Q_i^{v-1}. So we define $F_j^v=Q_j^v\setminus\Omega^{v+1}$, which are pairwise disjoint over all v and j. And since

$$|\Omega^{v+1}\cap Q_j^v|\le\frac{s}{1-t}|Q_j^v|,$$

we know that

$$|F_j^v|\ge\Big(1-\frac{s}{1-t}\Big)|Q_j^v|=c_{s,t}|Q_j^v|.$$

We can then compute

$$\begin{aligned}\inf_{y\in Q_j^v}M^\sharp_{0,s,\widehat{Q_j^v}}f(y)\int_{Q_j^v}w(x)dx &< c_{s,t}\frac{|F_j^v|}{|Q_j^v|}\inf_{y\in Q_j^v}M^\sharp_{0,s,Q_0}f(y)\int_{Q_j^v}w(x)dx\\ &= c_{s,t}\Big(\inf_{y\in Q_j^v}M^\sharp_{0,s,Q_0}f(y)\Big)^\delta|F_j^v|\frac{1}{|Q_j^v|}\int_{Q_j^v}\Big(M^\sharp_{0,s,Q_0}f(x)\Big)^{1-\delta}w(x)dx\\ &\le c_{s,t}\left(\int_{F_j^v}\Big(M^\sharp_{0,s,Q_0}f(x)\Big)^\delta dx\right)\inf_{y\in F_j^v}M\left(\Big(M^\sharp_{0,s,Q_0}f(\cdot)\Big)^{1-\delta}w(\cdot)\right)(y).\end{aligned}$$

Summing, we see that

$$\begin{aligned}&\sum_{v=1}^{\infty}\sum_{j\in I_2^v}\inf_{y\in Q_j^v}M^\sharp_{0,s,\widehat{Q_j^v}}f(y)\int_{Q_j^v}w(x)dx\\ &\qquad< c_{s,t}\int_{\mathbb{R}^n}\Big(M^\sharp_{0,s,Q_0}f(x)\Big)^\delta M\left(\Big(M^\sharp_{0,s,Q_0}f(\cdot)\Big)^{1-\delta}w(\cdot)\right)(x)dx.\end{aligned}$$

Similarly for $\sum_{v=1}^\infty\sum_{j\in I_2^v}\inf_{y\in Q_j^v}M^\sharp_{0,s,Q_j^v}f(y)$. □

Lerner's application of Theorem 2 is to Calderón-Zygmund singular integral operators via the inequality $M^\sharp_{0,s}(Tf)(x)\le cMf(x)$ [4]. Similar results follow for multipliers using inequalities derived in [10] (e.g., p.171).

A different estimate is needed in order for a local median oscillation decomposition to be applicable in Lerner's proof of the A_2 conjecture [7]. This new estimate leads to a similar local median oscillation decomposition, which we develop below.

Theorem 3. *Let f be a measurable function on a fixed cube $Q_0 \subset \mathbb{R}^n$, $0 < s < 1/2$, and $1/2 \leq t < 1-s$. Then there exists a (possibly empty) collection of subcubes $Q_j^v \in \mathcal{D}(Q_0)$ and a family of collections $\{I_2^v\}_v$ such that*

(i) *For a.e. $x \in Q_0$,*

$$|f(x) - m_f(t, Q_0)| \leq$$

$$8M^{\sharp}_{0,s,Q_0} f(x) + \sum_{v=1}^{\infty} \sum_j \left(m_{|f - m_f(t, \widehat{Q_j^v})|} \left(1 - (1-t)/2^n, \widehat{Q_j^v}\right) + m_{|f - m_f(t, Q_j^v)|}(t, Q_j^v) \right) \mathbb{1}_{Q_j^v}(x).$$

(ii) *For fixed v, $\{Q_j^v\}$ are pairwise disjoint families.*

(iii) *If $\Omega^v = \bigcup_j Q_j^v$, then $\Omega^{v+1} \subset \Omega^v$.*

(iv) *For all j, $|\Omega^{v+1} \cap Q_j^v| \leq \frac{s}{1-t} |Q_j^v|$.*

Proof. We follow the proof of Theorem 1 in form, with only a few definitional changes. First note that for any cube Q,

$$m_{|f - m_f(t,Q)|}(t, Q) \leq 4 \inf_{y \in Q} M^{\sharp}_{0,s,Q} f(y). \tag{18}$$

To see this, from (1.6), (1.7), (1.5), and (4.3) in [8],

$$\begin{aligned} m_{|f - m_f(t,Q)|}(t, Q) &\leq m_{|f - m_f(1-s,Q)| + |m_f(1-s,Q) - m_f(t,Q)|}(t, Q) \\ &\leq 2m_{|f - m_f(1-s,Q)|}(t, Q) \leq 2m_{|f - m_f(1-s,Q)|}(1-s, Q) \\ &\leq 4 \inf_{y \in Q} M^{\sharp}_{0,s,Q} f(y). \end{aligned}$$

We define $E^1 = \{x \in Q_0 : |f(x) - m_f(t, Q_0)| > m_{|f - m_f(t,Q_0)|}(t, Q_0)\}$. If $|E^1| = 0$, the decomposition halts, just as in Theorem 1. So we suppose $|E^1| > 0$. We then define

$$\Omega^1 = \{x \in Q_0 : m^{t,\Delta}_{Q_0}(f^0) > m_{|f - m_f(t,Q_0)|}(t, Q_0)\}.$$

Proceeding as above, we have that $\Omega^1 = \bigcup_j Q_j^1$ so that (as in (3))

$$|m_{f^0}(t, Q_j^1)| > m_{|f^0|}(t, Q_0), \; |m_f(t, \widehat{Q_j^1})| \leq m_{|f^0)|}(t, Q_0). \tag{19}$$

Furthermore, we also have that

$$\sum_j |Q_j^1| \leq \frac{s}{1-t} |Q_0|. \tag{20}$$

Before proceeding, observe that

$$m_f(t, Q) \leq m_f \left(1 - (1-t)/2^n, \widehat{Q}\right). \tag{21}$$

To see this,

$$\begin{aligned} |\{y \in \widehat{Q} : f(y) \geq m_f(t, Q)\}| &\geq |\{y \in Q : f(y) \geq m_f(t, Q)\}| \\ &\geq (1-t)|Q| = \frac{1-t}{2^n} \left|\widehat{Q}\right|, \end{aligned}$$

so taking complements in $\widehat{Q}$ we have

$$|\{y \in \widehat{Q} : f(y) < m_f(t, Q)\}| \leq \left(1 - \frac{1-t}{2^n}\right) |\widehat{Q}|.$$

Let $\alpha_j^1 = m_{f^0}(t, Q_j^1)$. By (19) and (21) we have

$$\begin{aligned} |\alpha_j^1| &\leq |m_{f^0}(t, Q_j^1) - m_{f^0}(t, \widehat{Q_j^1})| + |m_{f^0}(t, \widehat{Q_j^1})| \\ &\leq m_{|f^0 - m_{f^0}(t, \widehat{Q_j^1})|}(t, Q_j^1) + m_{|f^0|}(t, Q_0) \\ &\leq m_{|f - m_f(t, \widehat{Q_j^1})|}\left(1 - (1-t)/2^n, \widehat{Q_j^1}\right) + m_{|f^0|}(t, Q_0). \end{aligned} \tag{22}$$

This gives the first iteration of the local median oscillation decomposition of f when $|E^1| > 0$: for a.e. $x \in Q_0$, with $g^1 = f^0 \mathbb{1}_{Q_0 \setminus \Omega^1}$,

$$f^0(x) = g^1(x) + \sum_j \alpha_j^1 \mathbb{1}_{Q_j^1}(x) + \sum_j \left(f^0(x) - m_{f^0}(t, Q_j^1)\right) \mathbb{1}_{Q_j^1}(x).$$

Clearly by (18)

$$|g^1(x)| \leq m_{|f - m_f(t, Q_0)|}(t, Q_0) \leq 4 \inf_{y \in Q_0} M^\sharp_{0,s,Q_0} f(y) \leq 4 M^\sharp_{0,s,Q_0} f(x)$$

a.e. on $Q_0 \setminus \Omega^1$.

Proceeding as above, we assume that $\{Q_j^{k-1}\}$ are the dyadic cubes corresponding to the $(k-1)$st generation of subcubes of Q_0 obtained as above. Let

$$f_j^{k-1} = \left(f - m_f(t, Q_j^{k-1})\right) \mathbb{1}_{Q_j^{k-1}}$$

and

$$E_j^k = \{x \in Q_j^{k-1} : f_j^{k-1}(x) > m_{|f_j^{k-1}|}(t, Q_j^{k-1})\}.$$

If $|E_j^k| = 0$, we write

$$s_j^k = f_j^{k-1},$$

so that by (18), for a.e. $x \in Q_j^{k-1}$,

$$|s_j^k(x)| \leq m_{|f_j^{k-1}|}(t, Q_j^{k-1}) \leq 4 \inf_{y \in Q_j^{k-1}} M^\sharp_{0,s,Q_j^{k-1}} f(y) \leq 4 M^\sharp_{0,s,Q_0} f(x). \tag{23}$$

If $|E_j^k| > 0$, we define

$$\Omega_j^k = \{x \in Q_j^{k-1} : m^{t,\Delta}_{Q_j^{k-1}} f_j^{k-1}(x) > m_{|f_j^{k-1}|}(t, Q_j^{k-1})\}.$$

Then $|\Omega_j^k| > 0$ and

$$\Omega_j^k = \bigcup_i Q_i^k$$

where the Q_i^k's are nonoverlapping maximal dyadic subcubes of Q_j^{k-1} such that

$$|m_{f_j^{k-1}}(t,Q_i^k)| > m_{|f_j^{k-1}|}(t,Q_j^{k-1}),\ |m_{f_j^{k-1}}(t,\widehat{Q_i^k})| \le m_{|f_j^{k-1}|}(t,Q_j^{k-1}). \tag{24}$$

Then define

$$\Omega^k = \bigcup_j \Omega_j^k.$$

Let $\alpha_i^{k,j} = m_{f_j^{k-1}}(t,Q_i^k)$, and just as above

$$|\alpha_i^{k,j}| \le m_{|f-m_f(t,\widehat{Q_i^k})|}\left(1-(1-t)/2^n,\widehat{Q_i^k}\right) + m_{|f_j^{k-1}|}(t,Q_j^{k-1}). \tag{25}$$

We then have for a.e. $x \in Q_j^{k-1}$,

$$f_j^{k-1}(x) = g_j^k(x) + \sum_i \alpha_i^{k,j}\mathbb{1}_{Q_i^k}(x) + \sum_i \left(f(x) - m_f(t,Q_i^k)\right)\mathbb{1}_{Q_i^k}(x),$$

where, by (24) and (18), $g_j^k = f_j^{k-1}\mathbb{1}_{Q_j^{k-1}\setminus\Omega_j^k}$ satisfies

$$|g_j^k(x)| \le m_{|f_j^{k-1}|}(t,Q_j^{k-1}) \le 4\inf_{y\in Q_j^{k-1}} M^\sharp_{0,s,Q_j^{k-1}}f(y) \le 4M^\sharp_{0,s,Q_0}f(x) \tag{26}$$

for a.e. $x \in Q_j^{k-1}\setminus\Omega_j^k$. As in (13) we have that

$$|\Omega^k| \le \left(\frac{s}{1-t}\right)^k |Q_0|. \tag{27}$$

Furthermore, we have the kth iteration of the local median oscillation decomposition of f: for a.e. $x \in Q_0$,

$$f(x) - m_f(t,Q_0) = \sum_{v=1}^{k}\left(\sum_{j\in I_1^v} s_j^v + \sum_{j\in I_2^v} g_j^v\right) + \sum_{v=1}^{k}\sum_{j\in I_2^v}\sum_{i\in J_j^v}\alpha_i^{v,j}\mathbb{1}_{Q_i^v}(x) + \psi^k(x),$$

where

$$\psi^k = \sum_{j\in I_2^k}\sum_{i\in J_j^k}\left(f - m_f(t,Q_i^k)\right)\mathbb{1}_{Q_i^k}.$$

By (27), $\psi^k \to 0$ a.e. in Q_0 so that

$$\begin{aligned} f(x) - m_f(t,Q_0) &= \sum_{v=1}^{\infty}\left(\sum_{j\in I_1^v} s_j^v + \sum_{j\in I_2^v} g_j^v\right) + \sum_{v=1}^{\infty}\sum_{j\in I_2^v}\sum_{i\in J_j^v}\alpha_i^{v,j}\mathbb{1}_{Q_i^v}(x) \\ &= S_1(x) + S_2(x), \end{aligned}$$

say. As for S_1, by the same reasoning as in Theorem 1, (23) and (26) give that for a.e. $x \in Q_0$,

$$\begin{aligned} \left|\sum_{j\in I_1^v} s_j^v + \sum_{j\in I_2^v} g_j^v\right| &\le \max\left\{\max_{j\in I_1^v}\left\|f_j^{v-1}\right\|_{L^\infty}, \max_{j\in I_2^v}\left\|f_j^{v-1}\right\|_{L^\infty}\right\} \\ &\le 4M^\sharp_{0,s,Q_0}f(x). \end{aligned} \tag{28}$$

Now for S_2 and the $v=1$ summand, by (22),

$$\left|\sum_j \alpha_j^1 \mathbb{1}_{Q_j^1}(x)\right| \le \sum_j m_{|f-m_f(t,\widehat{Q_j^1})|}\left(1-(1-t)/2^n, \widehat{Q_j^1}\right)\mathbb{1}_{Q_j^1}(x) + m_{|f^0|}(t,Q_0)\mathbb{1}_{Q_0}. \tag{29}$$

As for the other terms, by (25) we have

$$\begin{aligned} \left|\sum_{v=2}^\infty \sum_{j\in I_2^v}\sum_{i\in J_j^v} \alpha_i^{v,j}\mathbb{1}_{Q_i^v}(x)\right| \le{}& \sum_{v=2}^\infty \sum_{j\in I_2^v}\sum_{i\in J_j^v} m_{|f-m_f(t,\widehat{Q_i^v})|}\left(1-(1-t)/2^n, \widehat{Q_i^v}\right)\mathbb{1}_{Q_i^v}(x) \\ &+ \sum_{v=2}^\infty \sum_{j\in I_2^v} m_{|f_j^{v-1}|}(t, Q_j^{v-1})\mathbb{1}_{Q_j^{v-1}}(x). \end{aligned} \tag{30}$$

Combining (29) and (30), and by the same reasoning as in Theorem 1,

$$\begin{aligned} \left|\sum_{v=1}^\infty \sum_{j\in I_2^v}\sum_{i\in J_j^v} \alpha_i^{v,j}\mathbb{1}_{Q_i^v}(x)\right| &\le \sum_{v=1}^\infty \sum_{j\in I_2^v}\sum_{i\in J_j^v} m_{|f-m_f(t,\widehat{Q_i^v})|}\left(1-(1-t)/2^n, \widehat{Q_i^v}\right)\mathbb{1}_{Q_i^v}(x) \\ &\quad + \sum_{v=1}^\infty \sum_{j\in I_2^v} m_{|f-m_f(t,Q_j^{v-1})|}(t, Q_j^{v-1})\mathbb{1}_{Q_j^{v-1}}(x) \\ &\le \sum_{v=1}^\infty \sum_{j\in I_2^v}\left(m_{|f-m_f(t,\widehat{Q_j^v})|}\left(1-(1-t)/2^n, \widehat{Q_j^v}\right) + m_{|f-m_f(t,Q_j^v)|}(t,Q_j^v)\right)\mathbb{1}_{Q_j^v} \\ &\quad + m_{|f-m_f(t,Q_0)|}(t,Q_0). \end{aligned} \tag{31}$$

Thus (18), (28), and (31) together give

$$\begin{aligned} &|f-m_f(t,Q_0)| \le \\ &\quad 8M^\sharp_{0,s,Q_0}f(x) + \sum_{v=1}^\infty \sum_{j\in I_2^v}\left(m_{|f-m_f(t,\widehat{Q_j^v})|}\left(1-(1-t)/2^n, \widehat{Q_j^v}\right) + m_{|f-m_f(t,Q_j^v)|}(t,Q_j^v)\right)\mathbb{1}_{Q_j^v} \end{aligned}$$

for a.e. $x \in Q_0$.

□

References

[1] D. Cruz-Uribe, J. M. Martell, and C. Perez, *Sharp weighted estimates for approximating dyadic operators*, Electron. Res. Announc. Math. Sci. **17** (2010), 12–19.

[2] N. Fujii, *A condition for the two-weight norm inequality for singular integral operators*, Studia Math. **98** (1991), no. 3, 175–190.

[3] J.B. Garnett and P.W. Jones, *BMO from dyadic BMO*, Pacific J. Math. **99** (1982), no. 2, 351–371.

[4] B. Jawerth and A. Torchinsky, *Local sharp maximal functions*, J. Approx. Theory (1985), no. 3, 231–270.

[5] A. Lerner, *A pointwise estimate for the local sharp maximal function with applications to singular integrals*, Bull. Lond. Math. Soc. **42** (2010), no. 5, 843–856.

[6] ———, *A "Local Mean Oscillation" Decomposition and Some of its Applications*, u.math.biu.ac.il/ lernera/paseky.pdf (2012).

[7] ———, *A Simple Proof of the A_2 Conjecture*, arxiv.org/abs/1202.2824 (2012).

[8] J. Poelhuis and A. Torchinsky, *Medians, Continuity, and Oscillation*, arxiv.org/abs/1207.2710 (2012).

[9] J.-O. Strömberg, *Bounded mean oscillation with Orlicz norms and duality of Hardy spaces*, Indiana Univ. Math. J. **28** (1979), no. 3, 511–544.

[10] J.-O. Strömberg and A. Torchinsky, *Weighted Hardy spaces*, Lecture Notes in Mathematics, vol. 1381, Springer-Verlag, Berlin, 1989.